

Pan Pearl River Delta Physics Olympiad 2021
2021 年泛珠三角及中华名校物理奥林匹克邀请赛
Sponsored by Institute for Advanced Study, HKUST
香港科技大学高等研究院赞助

Simplified Chinese Part-1 (Total 4 Problems, 40 Points) 简体版卷-1 (共4题, 40分)
(9:30 am – 12:00 pm, 15th May 2021)

Please fill in your final answers to all problems on the answer sheet.

请在答题纸上填上各题的最后答案。

At the end of the competition, please submit the answer sheet only. Question papers and working sheets will **not** be collected.

比赛结束时，请只交回答题纸，题目纸和草稿纸将**不会**收回。

1. [10 points]

A uniform thin rigid rod of mass m is supported by two rapidly rotating rollers, whose axes are separated by a fixed distance a . The rod is initially placed at rest symmetrically as shown in Fig.1a.

质量為 m 的均匀细刚性杆由两个快速旋转的滚筒支撑，两个滚筒的轴线距离為 a 。杆最初如图 1a 所示对称放置。

(a) [5 points] Assume that the rollers rotate in opposite directions as shown in Fig.1a. The coefficient of kinetic friction between the rod and the rollers is μ . Solve for the displacement $x(t)$ of the center C of the rod from roller 1 assuming $x(0) = x_0$ and $\dot{x}(0) = 0$.

(a) [5 分] 假设滚筒以相反方向旋转，如图 1a 所示。杆和滚筒之间的动摩擦系数为 μ 。假设 $x(0) = x_0$ 和 $\dot{x}(0) = 0$ ，求杆中心 C 从滚筒 1 量度的位移 $x(t)$ 。

(b) [5 points] Now consider the case in which the direction of rotation of the rollers are reversed, as shown in Fig.1b. Find the displacement $x(t)$, again assuming $x(0) = x_0$ and $\dot{x}(0) = 0$.

(b) [5 分] 现在考虑使滚筒的旋转方向反向的情况，如图 1b 所示。假设 $x(0) = x_0$ 和 $\dot{x}(0) = 0$ ，求杆中心 C 从滚筒 1 量度的位移 $x(t)$ 。




Fig. 1a





Fig. 1b

2. [10 points]

A cylindrical tank filled with liquid is placed on the side of the table. A small hole is opened on the wall of the tank near the bottom. The liquid flows out horizontally through the small hole, and hits at point P on the floor. The height of the table is H , the area of the hole is $\frac{1}{k}$ (assume $k \gg 1$) of the area of the bottom of the tank, and the initial height of the liquid in the tank is h_0 . Find the velocity v at which the drop point P moves along the floor and the time T required for all liquid to flow out of the tank.

在桌边放着装有液体的圆柱形容器，容器壁靠近底部开有小孔，液体经小孔水平流出，液柱射在地板上的 P 点。桌面高度为 H ，孔的面积是容器底部面积的 $\frac{1}{k}$ (假设 $k \gg 1$)，原来容器中液体高 h_0 。求落点 P 沿地板移动的速率 v 及所有液体从容器中流出所需的时间 T 。



3. [10 points]


A uniformly magnetized iron sphere of magnetic moment $\vec{m} = m\hat{z}$ and radius R is suspended from the ceiling by an insulating thread. A total charge $Q > 0$ is uniformly distributed throughout the iron sphere. We use the Cartesian coordinate where the origin is located at the center of the sphere, xy -plane is the horizontal plane and z -axis is pointing upward.

磁矩为 $\vec{m} = m\hat{z}$ 且半径为 R 的均匀磁化铁球通过绝缘线悬挂在天花板上。总电荷 $Q > 0$ 均匀分布在整個铁球中。我們使用笛卡尔坐标，其原点位于球体的中心， xy -平面是水平面， z 轴指向上方。

The magnetic field at position \vec{r} due to a magnetic dipole \vec{m} at origin is given by

由位於原點的磁偶極子 \vec{m} 所產生的磁場，在位置 \vec{r} 处为

$$\vec{B}(\vec{r}) = \frac{\mu_0}{4\pi r^3} \left(\frac{3(\vec{m} \cdot \vec{r})\vec{r}}{r^2} - \vec{m} \right)$$



(a) [3 points] The Poynting vector $\vec{S}(\vec{r})$ at point \vec{r} is defined as

(a) [3分] 坡印廷矢量 $\vec{S}(\vec{r})$ 在点 \vec{r} 上的定义为

$$\vec{S}(\vec{r}) = \frac{1}{\mu_0} \vec{E} \times \vec{B},$$

where \vec{E} and \vec{B} are the electric and magnetic fields at point \vec{r} . Calculate the magnitude of the Poynting vector $\vec{S}(\vec{r})$ both inside and outside of the iron sphere.

其中 \vec{E} 和 \vec{B} 为点 \vec{r} 的电场和磁场。计算铁球内部和外部的坡印廷矢量 $\vec{S}(\vec{r})$ 的大小。

(b) [2 points] In fact, the system has a non-zero angular momentum due to the electromagnetic field. The angular momentum L depends on the size of the sphere R , the charge Q and the magnetic dipole moment m on the sphere, and the physical constant μ_0 . We shall write $L = K\mu_0^\alpha R^\beta Q^\gamma m^\eta$ where K is a dimensionless constant. Find α, β, γ and η using dimensional analysis.

(b) [2分] 实际上，由于电磁场的存在，系统的角动量非零。角动量 L 取决于铁球的半径 R ，球上的电荷 Q 和磁偶极矩 m 以及物理常数 μ_0 。我们将写成 $L = K\mu_0^\alpha R^\beta Q^\gamma m^\eta$ ，其中 K 是无量纲常数。使用量纲分析找 α, β, γ 和 η 。

(c) [3 points] To calculate the total angular momentum \vec{L} of the system, it is given that the angular momentum density [angular momentum of the EM field per unit volume] $\vec{l}(\vec{r})$ of the electromagnetic field at a point \vec{r} is

(c) [3分] 为了计算系统的总角动量 \vec{L} ，已知在点 \vec{r} 处的电磁场角动量密度 [电磁场在每单位体积的角动量] $\vec{l}(\vec{r})$ 是

$$\vec{l}(\vec{r}) = \frac{1}{c^2} \vec{r} \times \vec{S}$$

where c is the speed of light. Calculate the total angular momentum \vec{L} of the system.

其中 c 是光速。计算系统的总角动量 \vec{L} 。

(d) [2 points] Electrons are injected into the iron sphere along the z-axis. The total amount of the charge in the sphere will reduce and the sphere will rotate. Find the angular speed of the sphere after the injection of N electrons. Assume that the moment of inertia of the iron sphere is I and each electron has charge $-e$.

(d) [2分] 电子沿 z 轴注入铁球。球体中的总电荷量减少，并且球体将旋转。找出 N 粒电子注入后球的角速度。假设铁球的惯性矩为 I ，并且每个电子的电荷为 $-e$ 。

4. Air Convection in Atmosphere 大气内之空气对流

(a) [1 point] Consider a horizontal slab of air whose thickness (height) is dz . If this slab is at rest, the pressure holding it up from below must balance both the pressure from above and the weight of the slab. Use this fact to find an expression for $\frac{dP}{dz}$, the variation of pressure with altitude, in terms of the density ρ of air.

(a) [1分] 考虑一薄片的空气，其厚度(高度)是 dz 。当这薄片处于静止状态时，从下方施加于薄片的压强必须平衡于从上面施加于薄片的压强和薄片的自身重量。由此，找出压强随高度变化的表达式 $\frac{dP}{dz}$ ，答案以空气密度 ρ 来表示。

(b) [2 points] Assume that the air is an ideal gas with molar mass M and the temperature T of the atmosphere is independent of height. Then the atmospheric pressure at height z is given by $P(z) = P(0)e^{-\lambda z}$. Find λ .

(b) [2 分] 假设空气是摩尔质量 M 的理想气体，而且大气的温度 T 随高度无关。因此，在高度为 z 的大气压强可以由 $P(z) = P(0)e^{-\lambda z}$ 表示。求 λ 。

In practice, the atmospheric temperature depends on height. If the temperature gradient $\left|\frac{dT}{dz}\right|$ exceeds a certain critical value, convection will occur: warm, low-density air will rise, while cool, high-density air sinks. The decrease of pressure with altitude causes a rising air mass to expand adiabatically and thus to cool. The condition for convection to occur is that the rising air mass must remain warmer than the surrounding air despite this adiabatic cooling.

在实际情况下，大气的温度会随高度变化。当温度梯度 $\left|\frac{dT}{dz}\right|$ 超越一个临界值时，对流就会产生：低密度的热空气上升，高密度的冷空气则下降。随高度上升而下降的气压，使上升的空气团发生绝热膨胀，从而冷却。对流发生的条件是：上升中的气团纵然发生绝热冷却，仍须较周围的空气温暖。

(c) [2 points] Assume that the molar heat capacity of air at constant volume is $c_V = \frac{5}{2}R$. We can show that when air expands adiabatically, the temperature and pressure are related by the condition

$$\frac{dT}{dP} = a \frac{T}{P}$$

Find the constant a .

(c) [2 分] 假设空气的定容摩尔热容量为 $c_V = \frac{5}{2}R$ 。由此可证明，当空气绝热膨胀时，温度和压强满足以下条件

$$\frac{dT}{dP} = a \frac{T}{P}$$

求常数 a 。

(d) [3 points] Assume that $\frac{dT}{dz}$ is just at the critical value for convection to begin, so that the temperature drop due to adiabatic expansion of the convecting air mass is the same as the temperature gradient of the surrounding air. Find a formula for $\frac{dT}{dz}$ in this case.

(d) [3 分] 假设 $\frac{dT}{dz}$ 正处于对流开始发生的临界值，以致对流空气由绝热膨胀引起的温度下降，等于周边空气的温度梯度。在此情况下，求 $\frac{dT}{dz}$ 的公式。

(e) [2 points] Calculate numerically the critical temperature gradient in part (d). Express your answer in K/km.

Data: The molar mass of the air is $M = 0.029 \text{ kg}$, $g = 9.8 \text{ m/s}^2$, $R = 8.31 \text{ J/mol/K}$, $T = 300\text{K}$.

(e) [2 分] 计算(d)部的临界温度梯度之数值。答案以 K/km 为单位。数值： $M = 0.029 \text{ kg}$, $g = 9.8 \text{ m/s}^2$, $R = 8.31 \text{ J/mol/K}$, $T = 300\text{K}$ 。

~ End of Part 1 卷-1 完 ~